



US010719692B2

(12) **United States Patent**
Kalscheur et al.

(10) **Patent No.:** **US 10,719,692 B2**

(45) **Date of Patent:** **Jul. 21, 2020**

(54) **VEIN MATCHING FOR DIFFICULT BIOMETRIC AUTHENTICATION CASES**

(71) Applicant: **Apple Inc.**, Cupertino, CA (US)

(72) Inventors: **Micah P. Kalscheur**, Cupertino, CA (US); **Feng Tang**, Cupertino, CA (US)

(73) Assignee: **Apple Inc.**, Cupertino, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/900,905**

(22) Filed: **Feb. 21, 2018**

(65) **Prior Publication Data**

US 2019/0080153 A1 Mar. 14, 2019

Related U.S. Application Data

(60) Provisional application No. 62/556,405, filed on Sep. 9, 2017, provisional application No. 62/556,812, filed on Sep. 11, 2017.

(51) **Int. Cl.**

G06K 9/00 (2006.01)

G06F 16/583 (2019.01)

G06F 16/56 (2019.01)

G06F 21/32 (2013.01)

(52) **U.S. Cl.**

CPC **G06K 9/00288** (2013.01); **G06F 16/56** (2019.01); **G06F 16/583** (2019.01); **G06F 16/5838** (2019.01); **G06F 21/32** (2013.01); **G06K 9/00255** (2013.01); **G06K 9/00281** (2013.01); **G06K 9/00885** (2013.01); **G06K 9/00892** (2013.01); **G06K 2009/00932** (2013.01)

(58) **Field of Classification Search**

CPC G06F 21/32; G06F 21/44; G06F 21/31; H04W 12/06

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,587,318 B2 9/2009 Seshadri
8,150,142 B2 4/2012 Freedman et al.
8,384,997 B2 2/2013 Shpunt et al.
8,635,066 B2 1/2014 Morrison
8,749,796 B2 6/2014 Pesach et al.
8,913,839 B2 12/2014 Ricanek, Jr. et al.

(Continued)

FOREIGN PATENT DOCUMENTS

WO 2016127766 A1 8/2016

OTHER PUBLICATIONS

ISR/WO, PCT/US2018/048791, dated Nov. 15, 2018, 17 pages.

(Continued)

Primary Examiner — Vincent Rudolph

Assistant Examiner — Raphael Schwartz

(74) *Attorney, Agent, or Firm* — Kowert, Hood, Munyon, Rankin & Goetzel, P.C.; Gareth M. Sampson

(57)

ABSTRACT

Subepidermal imaging of a face may be used to assess subepidermal features such as blood vessels (e.g., veins) when the device is attempting to authenticate a user in a facial recognition authentication process. Assessment of the subepidermal features may be used to distinguish between users that have closely related facial features (e.g., siblings or twins) in situations where the facial recognition authentication process has less certainty in a decision about recognition of the user's face as an authorized user.

20 Claims, 7 Drawing Sheets

